arXiv:1201.4187v1 [math.GT] 19 Jan 2012

Finite knot surgeries and Heegaard Floer homology

Margaret I. Doig

January 23, 2012

Abstract

It is well known that any 3-manifold can be obtained by Dehn surgery on a link but not which ones can be obtained from a knot or which knots can produce them. We investigate these two questions for elliptic Seifert fibered spaces (other than lens spaces) using the Heegaard Floer *correction terms* associated to a 3-manifold Y and its torsion Spin^c structures. If $H_1(Y)$ is finite and small, the correction terms completely identify the manifolds which can be realized as knot surgeries and the knots which realize them; even when $H_1(Y)$ is larger (but still finite), the invariants help identify the manifolds and place restrictions on the knots.

1 Introduction

In the 1960s, Wallace [Wal63] and Lickorish [Lic62] showed that any oriented 3-manifold can be constructed by Dehn surgery on a link in S^3. Soon after, Moser asked which manifolds can be constructed by surgery on a knot [Mos71]. One may also ask which knots give each manifold. We begin to answer these two questions for elliptic (or spherical) manifolds other than lens spaces (those with finite but non-cyclic fundamental group). We can structure the approach based on two classifications. By Thurston [Thu82], a knot is exactly one of the following: a torus knot (its complement is Seifert fibered and contains no essential torus), a hyperbolic knot (its complement admits a complete hyperbolic metric of finite volume), or a satellite knot (its complement contains an essential torus).

We know that S^3 only comes from trivial surgeries (see Moser [Mos71] for torus knots, Gordon and Luecke [GL89] for hyperbolic knots, and Gordon [Gor83a] for satellite knots), and $S^1 \times S^2$ arises only from 0-surgery on the unknot (see Gabai [Gab87]). On the other hand, lens spaces can come from the unknot and torus knots [Mos71] but may also arise from integral surgery [CGLS88] on some hyperbolic knots. Berge [Ber] suggested a list of such knots, the *primitive/primitive* knots, and Gordon conjectured that the list was complete in Problem 1.78 [Kir78a]. Ozsváth and Szabó [OS05] gave a necessary condition on the Alexander polynomial of a knot with a lens space surgery and verified Berge's list up to $p \leq 1500$, and Greene [Gre] verified that any lens space which is surgery on a non-]trivial knot is achieved by some knot on the list (he did not verify that all knots giving lens spaces are on the list). Dean [Dea03] attempted to extend these results from lens spaces to small Seifert fibered spaces. All of Dean's knots do have some fibered surgeries, although some infinite families of them have toroidal surgeries, too [EM02]. Other hyperbolic knots also admit small Seifert fibered surgeries, but the resulting manifolds can all be obtained by Dean's surgeries as well [DMM09, MMM06].

We address a subset of this case, elliptic (or spherical) manifolds other than the lens spaces, i.e., Seifert fibered manifolds with finite but non-cyclic fundamental group. We prove results such as Theorem 1, that the Poincaré homology sphere only comes from +1-surgery on the right-handed trefoil. Moser [Mos71] classified surgery on torus knots, showing that there are some finite and non-cyclic examples. Boyer and Zhang [BZ96] showed that any hyperbolic knot has at most six finite surgeries of distance five (if r_1 and r_2 are surgery coefficients, the distance $\Delta(r_1, r_2)$ is the minimal geometric intersection number of curves representing r_1 and r_2 on $\partial N(K)$, the boundary of a neighborhood of K). They also showed that a satellite knot with a finite surgery must be a cabled torus knot, and these knots were classified by Bleiler and Hodgson [BH92] with the help of Gordon's surgery transformation equation [Gor83b]; the transformation equation also tells us that any finite surgery on a satellite knot is also a finite surgery on a torus or hyperbolic knot. Beginning with manifolds of small first homology, we discuss some of these surgeries explicitly in Section 2.

Elliptic spaces fall into a group of manifolds called *L-spaces* whose Heegaard Floer homology is particularly simple. If an L-space is given by p/q surgery on a knot K in S^3, which we write $S^3_{p/q}(D)$, then the surgery coefficient and knot genus obey the inequality $p/q \geq 2g(K) - 1$, and the knot is fibered with one of a very small set of Alexander polynomials.

Since any elliptic space Y is a rational homology sphere, we may use an invariant called the *correction term* or *d-invariant* $d(Y, \mathfrak{t})$ for Y and an associated Spin^c structure, which are described in Section 3. For elliptic manifolds and many other Seifert fibered spaces, $d(Y, \mathfrak{t})$ may be calculated directly using a plumbing diagram. On the other hand, if Y is an L-space and a knot surgery, $d(Y, \mathfrak{t})$ can be derived from the Alexander polynomial and surgery coefficient.

By comparing the correction terms of an elliptic space to those of all the possible knot surgeries which may give it, we describe in Section 4 all of the surgeries of the elliptic spaces (other than lens spaces) with "small" first homology. For example, as conjectured by the remark after Problem 3.6(D) in [Kir78a] and Property I defined by Zhang [Zha93], the Poincaré homology sphere has a unique surgery construction:

Theorem 1. *The Poincaré homology sphere is* $+1$*-surgery on the right-handed trefoil (or, reversing orientation,* -1*-surgery on the left-handed trefoil)*[1]

$$\Sigma(2,3,5) = S^3_1(T_{2,3})$$

and no other surgery on any knot.

This result may also be seen from work of Ozsváth-Szabó, who calculate in Theorem 1.6 of [OS04d] that any knot giving the Poincaré homology sphere must have the same Alexander polynomial as the trefoil, and in [OS05] that it must be fibered.

Similarly,

Theorem 2. .

a. *The unique elliptic manifold (up to orientation other than lens spaces) with* $H_1(Y) = \mathbb{Z}_2$ *is*

$$S^3_2(T_{2,3})$$

and no other surgery on any knot.

---

[1] Throughout this paper, we suppress the choice of orientation and only mention positive surgeries.

b. The unique elliptic manifold (up to orientation other than lens spaces) with $H_1(Y) = \mathbb{Z}_3$ *is*

$$S^3_3(T_{2,3})$$

and no other surgery on any knot.

c. Of the infinite family of elliptic manifolds with $H_1(Y) = \mathbb{Z}_4$*, only one (up to orientation) is surgery on a knot in* S^3*, and it isxt*

$$S^3_4(T_{2,3})$$

and no other surgery on any knot.

When $H_1(Y)$ has more than four elements, the manifold may come from surgery on a knot of higher genus, which allows more unusual surgeries. (NB: There are no Seifert fibered manifolds with $H_1(Y) = \mathbb{Z}_5$ or $H_1(Y) = \mathbb{Z}_6$.)

Theorem 3. .

a. The unique elliptic manifold with $H_1(Y) = \mathbb{Z}_7$ *is*

$$S^3_{7/2}(T_{2,3}) \text{ and } S^3_7(K)$$

where K *is* $T_{2,5}$ *or a hyperbolic fibered knot with the same Alexander polynomial,* $\Delta_K(T) = T^2 - T + 1 - T^{-1} + T^{-2}$*.*

b. The infinite family of elliptic manifolds with $H_1(Y) = \mathbb{Z}_8$ *has only two members which are knot surgeries,*

$$S^3_8(T_{2,3}),$$

which is no other surgery on a knot, and

$$S^3_8(K),$$

where K *is* $T_{2,5}$ *or a hyperbolic fibered knot with the same Alexander polynomial.*

We summarize the results in Table 1 for all icosahedral, octahedral, and tetrahedral manifolds of $|H_1(Y)| \leq 10$ and all dihedral manifolds of $|H_1(Y)| \leq 32$.

The first presentation of this work may be found in [Doi10]. Special thanks go to Zoltan Szabó, who suggested and directed this project with a great deal of kindness and patience, and to Chuck Livingston, Paul Kirk, and Dave Gabai, who listened and contributed many helpful suggestions.



Figure 1: Kirby moves describing surgery on the right-handed trefoil.

2 Seifert fibered spaces as knot surgeries

A *p/q-[Dehn] surgery* on a knot K embedded in S^3 is an operation which removes an open neighborhood $N(K)$ homeomorphic to a solid torus and replaces it, identifying a meridian of the solid torus with $p\mu + q\lambda$ in the knot complement (where the meridian μ bounds a disk in $N(K)$ and the longitude λ is null-homologous in $H_1(S^3 - intN(K))$. Any closed, oriented 3-manifold Y is surgery on some link in S^3 [Lic62, Wal63]. A surgery diagram can be manipulated by the methods of Kirby calculus [Kir78b], which alter the diagram but not the underlying manifold: isotopy by surgery diagrams; stabilizing or destabilizing the manifold by addition or subtraction of a ±1-framed unknot which can be separated from the rest of the link; and handlesliding one link component over another, replacing L_2 with the band sum of L_1 and L_2. For the last, if n_i is the framing on L_i, then n_2 becomes $n_1 + n_2 + 2lk(L_1, L_2)$. See, e.g., Gompf and Stipsicz [GS99]. For example, p/q-surgery on the trefoil can be manipulated as in Figure 1.

Seifert fibered spaces. Seifert fibered spaces were originally defined by Seifert in 1932 ([Sei33], translated by W. Heil in [ST80]). Scott gives a more modern presentation with a slightly expanded definition in [Sco83].

A trivial solid torus may be given the product fibration. A *fibered solid torus* (or *fibered solid Klein bottle*) is a torus (or Klein bottle) which is finitely covered by the trivial fibered torus. A *Seifert fibered space* is a manifold fibered by circles so that any fiber has a neighborhood which is fiber isomorphic to a fibered solid torus or Klein bottle. Any exceptional (non-regular) fiber can be eliminated by some Dehn surgery, and the class of such surgery coefficients is referred to as the fiber's *framing.* A Seifert fibered space itself can be thought of as a fiber bundle over the orbifold obtained by compressing each fiber to a point.

For our purposes, we will need only Seifert fibered spaces whose base orbifolds are S^2. Construct such a space by choosing a line bundle ζ over S^2 and surgering over fibers with framings $-b_i/a_i$ (the negative sign is for historical reasons). It can be described as surgery on a link in S^3 whose components have framing $\{b = c_1(\zeta), -b_1/a_1, \ldots, -b_r/a_r\}$. Seifert identified such a manifold with an n-tuple (together with information about the base orbifold which we will exclude).

$$\left(b; \frac{a_1}{b_1}, \ldots, \frac{a_r}{b_r}\right)$$

For example, the p/q-surgery on the trefoil in Figure 1 is $\left(-1; \frac{1}{2}, \frac{1}{3}, \frac{q}{6q-p}\right)$. The choice of framings is not unique since a few handleslides will convert

$$\left(-b; \frac{a_1}{b_1}\right) \cong \left(-b-1; \frac{a_1}{b_1}+1\right)$$

Additionally, reversing the orientation as in [OS03a] shows

$$\left(-1; \frac{1}{2}, \frac{a_2}{b_2}, \frac{a_3}{b_3}\right) \cong -\left(-2; 1-\frac{1}{2}, 1-\frac{a_2}{b_2}, 1-\frac{a_3}{b_3}\right).$$

By geometrization [Pera, Perb, Perc], the manifolds with finite fundamental group are all Seifert fibered. They fall into five classes, those with cyclic π_1 and those with finite π_1 based on the four isometries of a sphere:

Theorem 4 (Seifert [Sei33]). *If Y^3 is closed, oriented, and Seifert-fibered with finite but non-cyclic fundamental group, then it has base orbifold S^2 and is one of:*

1. *Type I, icosahedral:* $\left(b; \frac{1}{2}, \frac{a_2}{3}, \frac{a_3}{5}\right)$ *with* $H_1(Y) = \mathbb{Z}_m$ *and* $(m, 30) = 1$.
2. *Type O, octahedral:* $\left(b; \frac{1}{2}, \frac{a_2}{3}, \frac{a_3}{4}\right)$ *with* $H_1(Y) = \mathbb{Z}_{2m}$ *and* $(m, 24) = 1$.
3. *Type T, tetrahedral:* $\left(b; \frac{1}{2}, \frac{a_2}{3}, \frac{a_3}{3}\right)$ *with* $H_1(Y) = \mathbb{Z}_{3m}$ *and* $(m, 18) = 1$.
4. *Type D, dihedral:* $\left(b; \frac{1}{2}, \frac{1}{2}, \frac{a_3}{b_3}\right)$ *with* $H_1(Y) = \mathbb{Z}_{4m}$ *and* $(m, 4b_3) = 1$ *(if* b_3 *is even) or* $H_1(Y) = \mathbb{Z}_2 \times \mathbb{Z}_{2m}$ *(if* b_3 *is odd).*

with coefficients chosen always so that $(a_i, b_i) = 1$.

Specifically, $H_1(Y) = \mathbb{Z}_p$ and $p = b_1 b_2 b_3 \left(-b + \frac{a_1}{b_1} + \frac{a_2}{b_2} + \frac{a_3}{b_3}\right)$. In fact, if an integer p meets the requirements for one of the four types, there is a unique Seifert fibered space with $H_1(Y) = \mathbb{Z}_p$ of types I, O, or T, or a unique infinite family of type D. Each space has a unique description (possibly with reversed orientation) with $b = -1$; $0 < a_1 < b_1$, $0 < a_2 < b_2$, and $0 < a_3$; and $(a_i, b_i) = 1$.

Using Theorem 4, we include in Table 1 all the type I, O, or T manifolds with $|H_1(Y)| \leq 10$ and all type D with $|H_1(Y)| \leq 32$.

Finite surgeries. Many of the elliptic manifolds can be realized as torus knot surgeries.

Theorem 5 (Moser [Mos71])**.**

$$S^3_{p/q}(T_{r,s}) = \begin{cases} L_{r,s} \# L_{s,r} & if\ p/q = rs \\ \left(b; \frac{a_1}{r}, \frac{a_2}{s}, \frac{a_3}{|rsq-p|}\right) & if\ p/q \neq rs\ for\ some\ a_1, a_2, a_3 \\ L_{p,rsq} & if\ p/q = rs \pm 1/q \end{cases}$$

Corollary 6. *The torus knots which give elliptic spaces (other than lens spaces) are:*

$$S^3_{p/q}(T_{2,s}) = \left(-1; \frac{1}{2}, \frac{(s-1)/2}{s}, \frac{q}{2sq-p}\right) = -\left(-1; \frac{1}{2}, \frac{(s+1)/2}{s}, \frac{q}{p-2sq}\right)$$
$$S^3_{p/q}(T_{3,4}) = \left(-1; \frac{2}{3}, \frac{1}{4}, \frac{q}{12q-p}\right) = -\left(-1; \frac{1}{3}, \frac{3}{4}, \frac{q}{p-12q}\right)$$
$$S^3_{p/q}(T_{3,5}) = \left(-1; \frac{2}{3}, \frac{2}{5}, \frac{q}{15q-p}\right) = -\left(-1; \frac{1}{3}, \frac{3}{5}, \frac{q}{p-15q}\right)$$

$H_1(Y)$	*type*	*Seifert description*	*knot surgeries*
1	I	$(-1; \frac{1}{2}, \frac{1}{3}, \frac{1}{5})$	$S^3_1(T_{2,3})$
$\mathbb{Z}_2$	O	$(-1; \frac{1}{2}, \frac{1}{3}, \frac{1}{4})$	$S^3_2(T_{2,3})$
$\mathbb{Z}_3$	T	$(-1; \frac{1}{2}, \frac{1}{3}, \frac{1}{3})$	$S^3_3(T_{2,3})$
$\mathbb{Z}_4$	D	$(-1; \frac{1}{2}, \frac{1}{2}, \frac{1}{4})$	$S^3_4(T_{2,3})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{1}{n})$	*none if* $n > 4$
$\mathbb{Z}_7$	I	$(-1; \frac{1}{2}, \frac{1}{3}, \frac{2}{5})$	$S^3_{7/2}(T_{2,3}) = S^3_7(T_{2,5})$
$\mathbb{Z}_8$	D	$(-1; \frac{1}{2}, \frac{1}{2}, \frac{2}{3})$	$-S^3_8(T_{2,3})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{2}{5})$	$S^3_8(T_{2,5})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{2}{n})$	*none if* $n > 5$
$\mathbb{Z}_9$	T	$(-1; \frac{1}{2}, \frac{1}{3}, \frac{2}{3})$	$S^3_{9/2}(T_{2,3})$
$\mathbb{Z}_{10}$	O	$(-1; \frac{1}{2}, \frac{2}{3}, \frac{1}{4})$	$-S^3_{10}(T_{2,3}) = S^3_{10}(T_{3,4})$
$\mathbb{Z}_{12}$	D	$(-1; \frac{1}{2}, \frac{1}{2}, \frac{3}{5})$	$-S^3_{12}(T_{2,5})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{3}{7})$	$S^3_{12}(T_{2,7})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{3}{n})$	*none if* $n > 7$
$\mathbb{Z}_{16}$	D	$(-1; \frac{1}{2}, \frac{1}{2}, \frac{4}{3})$	$S^3_{16/3}(T_{2,3})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{4}{7})$	$-S^3_{16}(T_{2,7})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{4}{9})$	$S^3_{16}(T_{2,9})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{4}{n})$	*none if* $n \neq 3, 7, 9$
$\mathbb{Z}_{20}$	D	$(-1; \frac{1}{2}, \frac{1}{2}, \frac{5}{3})$	$-S^3_{20/3}(T_{2,3})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{5}{9})$	$-S^3_{20}(T_{2,9})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{5}{11})$	$S^3_{20}(T_{2,11})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{5}{n})$	*none if* $n \neq 3, 9, 11$
$\mathbb{Z}_{24}$	D	$(-1; \frac{1}{2}, \frac{1}{2}, \frac{6}{11})$	$-S^3_{24}(T_{2,11})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{6}{13})$	$S^3_{24}(T_{2,13})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{6}{n})$	*none if* $n \neq 11, 13$
$\mathbb{Z}_{28}$	D	$(-1; \frac{1}{2}, \frac{1}{2}, \frac{7}{3})$	$S^3_{28/5}(T_{2,3})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{7}{13})$	$-S^3_{28}(T_{2,13})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{7}{15})$	$S^3_{28}(T_{2,15})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{7}{n})$	*none if* $n \neq 3, 13, 15$
$\mathbb{Z}_{32}$	D	$(-1; \frac{1}{2}, \frac{1}{2}, \frac{8}{3})$	$-S^3_{32/5}(T_{2,3})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{8}{15})$	$-S^3_{32}(T_{2,15})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{8}{17})$	$S^3_{32}(T_{2,17})$
		$(-1; \frac{1}{2}, \frac{1}{2}, \frac{8}{n})$	*none if* $n \neq 3, 15, 17$

Table 1: Seifert fibered spaces of types I, O, and T with $H_1(Y) \leq 10$ and type D with $|H_1(Y)| \leq 32$ and their knot surgery constructions.

For example, any icosahedral manifold comes from at least a trefoil surgery (because $(p, 30) = 1$ so $p \equiv \pm 5 \bmod 6$, so either $q = (p+5)/6$ or $(p-5)/6$ is an integer) and sometimes also $T_{2,5}$ or $T_{3,5}$. Similarly, any octahedral comes from $T_{2,3}$ and possibly $T_{3,4}$, any tetrahedral from $T_{2,3}$, and a dihedral may come from $T_{2,3}$ or $T_{2,s}$ if $s = (p/4 \pm 1)/2$.

Bleiler and Hodgson list finite surgeries on iterated torus knots [BH96] based on Gordon [Gor83a]. Boyer and Zhang prove that any finite filling of a hyperbolic knot is either integral or half-integral (which cannot happen for tetrahedral or dihedral manifolds); that no hyperbolic knot has both a cyclic and a finite (non-cyclic) filling; that no satellite knot other than a 1- or 2-iterated torus knot has a finite filling; and that there is a bound on the number of and distance between finite/cyclic fillings [BZ96, BZ01]. Eudave-Munoz and others describe some peculiar fillings that may give elliptic spaces [EM02].

3 The invariant $d(Y, \mathfrak{t})$

Heegaard Floer homology [OS04c, OS04b] is an example of a Floer homology which assigns a set of invariants (in our case, a graded abelian group over $\mathbb{Z}/2$) to a closed, connected, oriented 3-manifold using a Heegaard decomposition of the manifold. A Floer homology starts with a $2n$-dimensional symplectic manifold and two n-dimensional Lagrangian submanifolds which meet transversely. The chain complex is a free R-module (for $R = \mathbb{Z}_2$, $\mathbb{Z}$, etc.) whose generators come from intersection points of the Lagrangians and whose boundary map connects two generators if there is an embedded "disk" between them. Heegaard Floer homology defines the symplectic manifold and the Lagrangians using a Heegaard decomposition of a 3-manifold. The generators of its chain complex can be thought of as sets of points on the Heegaard surface and the boundary maps as domains in the surface.

Heegaard Floer assigns a set of invariants to Y and its Spin^c structures, the *correction terms* or *d-invariants* $d(Y, \mathfrak{t})$. The hat version $\widehat{HF}(Y)$ comes with a relative $\mathbb{Z}$-grading which lifts to an absolute $\mathbb{Q}$-grading for a rational homology sphere (see Theorem 7.1 of [OS06]); it is defined by requiring that $\widehat{HF}(S^3) \cong \mathbb{Z}$ is supported in degree 0 and that the inclusion map $\widehat{CF}(Y, \mathfrak{t}) \hookrightarrow CF^+(Y, \mathfrak{t})$ preserves degree. Then $d(Y, \mathfrak{t})$ is the minimal grading of any non-torsion class in $HF^+(Y, \mathfrak{t})$ coming from $HF^\infty(Y, \mathfrak{t})$ [OS03a]. (If Y is elliptic, each Spin^c structure corresponds to a torsion element of $H_1(Y, \mathbb{Z})$, all classes in $HF^+(Y)$ come from $HF^\infty(Y)$, and $d(Y, \mathfrak{t})$ is defined for all $\mathfrak{t}$.)

***L*-space surgeries.** An elliptic Seifert fibered space is an example of an L-space, the Heegaard Floer homology version of a lens space [OS03b]. $\widehat{HF}(Y)$ splits into $\oplus^{\mathfrak{t}}\widehat{HF}(Y, \mathfrak{t})$ over Spin^c structures (equivalence classes of non-zero vector fields $\mathfrak{t}$ isomorphic to $H^2(Y; \mathbb{Z})$ by $\mathfrak{t} \mapsto c_1(\mathfrak{t})$). Lens spaces have the nice property that each generator of $\widehat{HF}(L_{p,q})$ falls into a different torsion Spin^c structure ($\mathfrak{t} \in Spin^c(Y)$ is torsion if $PD(c_1(\mathfrak{t}) \in H_1(Y)$ is torsion). We will call any rational homology sphere with this property an *L-space.* Equivalently, $\widehat{HF}(Y, \mathfrak{t}) \cong \widehat{HF}(S^3)$.

Using the surgery exact sequences and absolute grading on $HF^+(Y)$, we can place some restrictions on which knots may have L-space surgeries. Normalize the Alexander polynomial so

$$\Delta_K(T) = a_0 + \sum_{i=1}^{n} a_i(T^i + T^{-i}).$$

Theorem 7 (Theorem 1.2 [OS04a]; Corollary 1.3 [OS05])**.** *If a knot $K \subset S^3$ admits an L-space surgery, then the non-zero coefficients of $\Delta_K(T)$ are alternating $+1$s and -1s.*

Ozsváth and Szabó [OS04a] showed that the knot Floer homology $\widehat{HFK}(K, i)$ is $\mathbb{Z}$ in the top grading $i = g(K)$ for any fibered knot, and Ni [Ni09] and, independently, Juhász [Juh06] showed the reverse; since $\Delta_K(T)$ is the graded Euler characteristic of $\widehat{HFK}(S^3, K)$, this means that

Corollary 8. *If a knot $K \subset S^3$ admits an L-space surgery, then K is fibered.*

Finally,

Theorem 9 ([OS, OS08])**.** *If K admits a positive L-space surgery, then $S^3_{p/q}(K)$ is an L-space if and only if*

$$\frac{p}{q} \geq 2g(K) - 1.$$

Calculating $d(Y,\mathfrak{t})$ of a knot surgery. If $S^3_{p/q}(K)$ is an L-space, then $\widehat{HF}(S^3_{p/q}(K))$ and its gradings can be calculated from $\Delta_K(T)$ and p/q:

Theorem 10. *If $0 < q < p$,*

a. Proposition 4.8, [OS03a]: for $0 \leq i < p+q$,

$$d(S^3_{p/q}(U), i) = -\left(\frac{pq - (2i+1-p-q)^2}{4pq}\right) - d(S^3_{q/r}(U), j)$$

where $r \equiv p \bmod q$ and $j \equiv i \bmod q$.

b. Theorem 1.2, [OS]: for $|i| \leq \frac{p}{2}$,

$$d(S^3_{p/q}(K), i) - d(S^3_{p/q}(U), i) = -2\sum_{j=1}^{\infty} j a_{c+j}$$

where $c = \left|\left\lfloor \frac{i}{q} \right\rfloor\right|$.

Calculating $d(Y,\mathfrak{t})$ of a Seifert fibered space. It is sometimes also possible to calculate the $d(Y,\mathfrak{t})$ using a negative definite plumbing diagram [OS03a, OS03b].

Any 3-manifold Y is the boundary of some oriented 4-manifold. For example, $p/1$-surgery on a knot in Y is cobordant to Y (attach a 2-handle to $Y \times [0,1]$ by gluing its boundary along the image of the knot in $Y \times \{1\}$ with framing p.) The same diagram describing Y as integral surgery on a link L can describe X by gluing 2-handles along L, which is called a *plumbing diagram* for X. Say L is a set of unknots which link at most once pairwise, so we can describe Y by drawing a graph G with a vertex for each component of L and an edge if two components are linked. For example, a surgery diagram for $Y = \left(-2; \frac{1}{2}, \frac{1}{2}, \frac{2}{3}\right)$ might be



(NB: $Y \cong -\left(-1; \frac{1}{2}, \frac{1}{2}, \frac{1}{3}\right)$; we have reversed the orientation for this calculation.) Label each vertex v with a weight $m(v)$ which is the framing of that component.

Similarly, the same graph describes a 4-manifold cobordism X bounded by Y. Each vertex represents the cocore of the corresponding handle, which is a generator of $H^2(X, \partial X; \mathbb{Z})$, and an edge exists if two such generators intersect. The weight of a vertex v gives the self-intersection number $\langle PD^{-1}(v), PD^{-1}(v)\rangle$ of the corresponding generator. In fact, the intersection form of X

$$Q_X : H^2(X, \partial X; \mathbb{Z}) \times H^2(X, \partial X; \mathbb{Z}) \to \mathbb{Z}$$
$$Q_X(\alpha, \beta) = \langle \alpha, \beta \rangle = \#(a \cap b)$$

can be read off from this description, which is the symmetric bilinear form assigning to $\langle \alpha, \beta\rangle$ the algebraic intersection number of surfaces representing their Poincaré duals a and b. Then the *characteristic vectors* or $Char(G)$ of Q_X are the $\alpha \in H^2(X, \partial X; \mathbb{Z})$ such that

$$\langle \alpha, \beta \rangle = \langle \beta, \beta \rangle \quad \forall \beta \in H^2(X, \partial X; \mathbb{Z}) \quad \mod 2.$$

According to [OS03b], $HF^+(Y)$ can be expressed by certain functions on the characteristic vectors of the intersection matrix. Let

$$\mathcal{T}_0^+ = \mathbb{Z}[U, U^{-1}]/U \cdot \mathbb{Z}[U]$$

as a $\mathbb{Z}[U]$-module with grading so that U^{-d} is homogeneous and supported in degree $2d$ (where $d > 0$). Consider the set $\mathbb{H}^+(G)$ of all homomorphisms

$$\phi : Char(G) \to \mathcal{T}_0^+$$

which preserve the adjunction relations

$$\begin{aligned} U^n \phi(\alpha + PD^{-1}[v]) &= \phi(\alpha) && \text{if } n \geq 0 \\ \phi(\alpha + PD^{-1}[v]) &= U^{-n}\phi(\alpha) && \text{if } n \leq 0 \end{aligned}$$

for $\alpha \in Char(G)$, any $v \in G$, and where $2n = \langle \alpha, PD^{-1}(v)\rangle + \langle PD^{-1}(v), PD^{-1}(v)\rangle$. Then $\mathbb{H}^+(G)$ splits over $\mathrm{Spin}^c(Y)$ and can be given a grading adapted from the one on $\mathcal{T}_0^+$: if $\phi(\alpha) \in \mathcal{T}_0^+$ is a homogeneous element and $\alpha \in Char(G)$ so that $\phi(\alpha) \neq 0$, then

$$\deg(\phi) = \deg\left(\phi(\alpha) \subset \mathcal{T}_0^+\right) - \frac{\alpha^2 + |G|}{4}$$

Let G be a negative definite weighted graph with at most two *bad vertices*, or vertices where $\deg(v) > |m(v)|$. Then

Theorem 11.

$$HF^+(-Y(G), \mathfrak{t}) \cong \mathbb{H}^+(G, \mathfrak{t})$$

and the grading on $\mathbb{H}^+(G)$ *is the absolute* $\mathbb{Q}$*-grading on* HF^+*.*

Corollary 12. *If* $Char_{\mathfrak{t}}(G)$ *is the set of characteristic vectors* $k = c_1(\mathfrak{t})$*,*

$$d(-Y(G), \mathfrak{t}) = \max_{\alpha \in Char_{\mathfrak{t}}(G)} \frac{\alpha^2 + |G|}{4}.$$

For a given $\mathfrak{t}$, the maximum is achieved among the finite set of vectors with

$$|\langle \alpha, PD^{-1}(v) \rangle| \leq |m(v)|$$

for all v. By [OS03b], these vectors are exactly the ones with representatives of the form $U^m \alpha$ for $m > 0$. For each of these classes, all the vectors obey $|\langle \alpha, PD^{-1}(v) \rangle| \leq |m(v)|$, and there are unique vectors α and β with

$$\begin{aligned} m(v) &\leq \langle \alpha, PD^{-1}(v) \rangle < -m(v) \\ m(v) &< \langle \beta, PD^{-1}(v) \rangle \leq -m(v) \end{aligned} \tag{1}$$

Any other vector in the class belongs to a [usually not unique] *full path* from α to β, meaning a sequence of vectors $(\alpha = \alpha_0, \alpha_1, \cdots, \alpha_n = \beta)$ where

$$\alpha_i = \alpha_{i-1} + 2PD[v_{i+1}] \qquad \text{and} \qquad \langle \alpha_i, PD^{-1}(v_{i+1}) \rangle = -m(v_{i+1}).$$

This property is unique to the classes without any $U^m \alpha$ for $m > 0$. Both α and β are independent of the particular path chosen, as long as $\langle \alpha_i, PD^{-1}(v_{i+1}) \rangle = -m(v_{i+1})$ at each step, and $\langle \alpha_i, \alpha_i \rangle$ is constant for each full path.

Corollary 13. *If* $\widehat{Char_{\mathfrak{t}}(G)}$ *is the subset of* $Char_{\mathfrak{t}}(G)$ *which are part of a full path obeying (1), then*

$$d(-Y(G), \mathfrak{t}) = \max_{\alpha \in \widehat{Char_{\mathfrak{t}}(G)}} \frac{\alpha^2 + |G|}{4}.$$

Example 14. For example, we may wish to calculate the correction terms for the manifolds $Y = \left(-1; \frac{1}{2}, \frac{1}{2}, \frac{1}{n}\right)$, which are listed in Table 2.

The trees for these manifolds are easier to manipulate if we reverse orientation: $-Y = -\left(-2; \frac{1}{2}, \frac{1}{2}, \frac{n-1}{n}\right)$, which is negative definite since $n-1 < n$. It has plumbing diagram



where there are $k = n-1$ vertices to the right of the central node and each unlabeled vertex has weight -2, i.e., $\langle v_i, v_i \rangle = -2$. [2]

Equivalently, the 4-manifold bounded by Y has intersection form

$$Q = \begin{bmatrix} -2 & 1 & 1 & 1 & 0 & \cdots & 0 & 0 \\ 1 & -2 & 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 0 & -2 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 0 & 0 & -2 & 1 & \cdots & 0 & 0 \\ 0 & 0 & 0 & 1 & -2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & -2 & 1 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 1 & -2 \end{bmatrix}$$

To evaluate a characteristic vector α, it may be easier to consider the vector $Q\alpha$; we will list the characteristic vectors by drawing the plumbing diagram and labeling it with the vector $Q\alpha$, i.e., the vertex v_i gets the label $e_i^T Q\alpha$ if $e_i = PD^{-1}(v_i)$. The characteristic vectors are:









---

[2] We will usually suppress a vertex's weight if it is -2, and we will number the vertices in order starting with the central node (which is v_1), then the left arm (v_2), followed by the bottom arm (v_3), and finally the right arm ($v_4, \cdots, v_{k+3}$).

Since $\langle \alpha, \alpha \rangle = \alpha^2 = \alpha^T Q \alpha = (Q\alpha)^T Q^{-1} (Q\alpha)$, it is straightforward to calculate

$$\begin{aligned} {\alpha_0}^2 &= 0 \\ {\alpha_1}^2 &= {\alpha_{-1}}^2 = -k-3 \\ {\alpha_2}^2 &= -4 \end{aligned}$$

so

$$d(-Y(G), \mathfrak{t}_i) = \frac{{\alpha_i}^2 + k + 3}{4} = \begin{cases} \frac{n+2}{4} & i = 0 \\ 0 & i = \pm 1 \\ \frac{n-2}{4} & i = 2 \end{cases}$$

4 $d(Y, \mathfrak{t})$ as a knot surgery obstruction

We will begin to identify the finite, non-cyclic surgeries by looking at all elliptic manifolds (other than lens spaces) of $|H_1(Y)| \leq 10$ and all dihedral manifolds of $|H_1(Y)| \leq 32$. We list all such manifolds which are knot surgeries, and we exhibit at least one knot that produces each surgery and say something about which other knots may do so. This latter question is one of *cosmetic surgery* (a non-trivial surgery that does not change the diffeomorphism type of the manifold; i.e., it changes the manifold in appearance but not in substance).

Proof of Theorem 1. Since p/q-surgery on a knot in an integral homology sphere gives $H_1(Y) = \mathbb{Z}_p$, assume the surgery coefficient is some rational number $1/q$. Since $S^3_{p/q}(K) = -S^3_{-p/q}(mK)$ where mK is the mirror of K, and since $d(Y, \mathfrak{t}) = -d(-Y, -\mathfrak{t})$, we will suppress further discussion of negative coefficients and assume $p/q \geq 0$.

By Theorem 4, the Poincaré homology sphere is

$$Y = \left(-1; \frac{1}{2}, \frac{1}{3}, \frac{1}{5} \right).$$

We ignore lens spaces, so $g(K) > 0$; also, this manifold is an L-space, so Theorem 9 restricts q:

$$\frac{1}{q} \geq 2g(K) - 1$$

p	k										
4	1	0	$\frac{n+2}{4}$	$\frac{n-2}{4}$							
8	1	$\frac{1}{4}$	$-\frac{1}{4}$	$\frac{n-4}{8}$	$\frac{n+4}{8}$						
12	1	$\frac{1}{6}$	$-\frac{1}{2}$	$\frac{n+4}{12}$	$\frac{n-8}{12}$	$\frac{n-4}{12}$	$\frac{n+8}{12}$				
12	2	$\frac{1}{2}$	$-\frac{1}{6}$	$\frac{n-4}{12}$	$\frac{n+8}{12}$	$\frac{n+4}{12}$	$\frac{n-8}{12}$				
16	1	0	$\frac{1}{4}$	$-\frac{3}{4}$	$\frac{n+2}{16}$	$\frac{n-14}{16}$	$\frac{n-6}{16}$	$\frac{n+10}{16}$			
16	3	0	$-\frac{1}{4}$	$\frac{3}{4}$	$\frac{n-2}{16}$	$\frac{n+14}{16}$	$\frac{n+6}{16}$	$\frac{n-10}{16}$			
20	1	-1	$\frac{1}{5}$	$-\frac{1}{5}$	$\frac{n-2}{20}$	$\frac{n-22}{20}$	$\frac{n+10}{20}$	$\frac{n-10}{20}$	$\frac{n+14}{20}$	$\frac{n-6}{20}$	
20	2	0	$\frac{2}{5}$	$-\frac{2}{5}$	$\frac{n-2}{20}$	$\frac{n+18}{20}$	$\frac{n-10}{20}$	$\frac{n+10}{20}$	$\frac{n-14}{20}$	$\frac{n+6}{20}$	
20	3	0	$\frac{2}{5}$	$-\frac{2}{5}$	$\frac{n+2}{20}$	$\frac{n-18}{20}$	$\frac{n+10}{20}$	$\frac{n-10}{20}$	$\frac{n+14}{20}$	$\frac{n-6}{20}$	
20	4	1	$\frac{1}{5}$	$-\frac{1}{5}$	$\frac{n+2}{20}$	$\frac{n+22}{20}$	$\frac{n-10}{20}$	$\frac{n+10}{20}$	$\frac{n-14}{20}$	$\frac{n+6}{20}$	
24	1	$\frac{1}{4}$	$\frac{1}{12}$	$-\frac{5}{12}$	$-\frac{5}{4}$	$\frac{n-10}{24}$	$\frac{n+14}{24}$	$\frac{n+6}{24}$	$\frac{n-18}{24}$	$\frac{n-34}{24}$	$\frac{n-10}{24}$
24	5	$-\frac{1}{4}$	$-\frac{1}{12}$	$\frac{5}{12}$	$\frac{5}{4}$	$\frac{n-10}{24}$	$\frac{n+14}{24}$	$\frac{n+6}{24}$	$\frac{n-18}{24}$	$\frac{n+30}{24}$	$\frac{n+6}{24}$
28	1	$\frac{3}{14}$	$-\frac{1}{14}$	$-\frac{9}{14}$	$-\frac{3}{2}$						
				$\frac{n-18}{28}$	$\frac{n-46}{28}$	$\frac{n+2}{28}$	$\frac{n-26}{28}$	$\frac{n+14}{28}$	$\frac{n-14}{28}$	$\frac{n+18}{28}$	$\frac{n-10}{28}$
28	2	$\frac{3}{14}$	$-\frac{1}{14}$	$-\frac{9}{14}$	$\frac{1}{2}$						
				$\frac{n+8}{28}$	$\frac{n-20}{28}$	$\frac{n-4}{28}$	$\frac{n+24}{28}$	$\frac{n-24}{28}$	$\frac{n+4}{28}$	$\frac{n-16}{28}$	$\frac{n+12}{28}$
28	3	$-\frac{3}{14}$	$\frac{1}{14}$	$\frac{9}{14}$	$-\frac{1}{2}$						
				$\frac{n+18}{28}$	$\frac{n-10}{28}$	$\frac{n+2}{28}$	$\frac{n+30}{28}$	$\frac{n-22}{28}$	$\frac{n+6}{28}$	$\frac{n-18}{28}$	$\frac{n+10}{28}$
28	4	$\frac{3}{14}$	$-\frac{1}{14}$	$-\frac{9}{14}$	$\frac{1}{2}$						
				$\frac{n+8}{28}$	$\frac{n-20}{28}$	$\frac{n-4}{28}$	$\frac{n-32}{28}$	$\frac{n+20}{28}$	$\frac{n-8}{28}$	$\frac{n+16}{28}$	$\frac{n-12}{28}$
28	5	$-\frac{3}{14}$	$\frac{1}{14}$	$\frac{9}{14}$	$-\frac{1}{2}$						
				$\frac{n+4}{28}$	$\frac{n-24}{28}$	$\frac{n-4}{28}$	$\frac{n+24}{28}$	$\frac{n+20}{28}$	$\frac{n-8}{28}$	$\frac{n-12}{28}$	$\frac{n+16}{28}$
28	6	$-\frac{3}{14}$	$\frac{1}{14}$	$\frac{9}{14}$	$\frac{3}{2}$						
				$\frac{n+14}{28}$	$\frac{n+42}{28}$	$\frac{n+6}{28}$	$\frac{n-22}{28}$	$\frac{n-6}{28}$	$\frac{n+22}{28}$	$\frac{n+10}{28}$	$\frac{n-18}{28}$
32	1	$\frac{1}{8}$	$\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{7}{8}$	$-\frac{7}{4}$					
				$\frac{n-26}{32}$	$\frac{n-58}{32}$	$\frac{n-2}{32}$	$\frac{n-34}{32}$	$\frac{n+14}{32}$	$\frac{n-18}{32}$	$\frac{n+22}{32}$	$\frac{n-10}{32}$
32	3	$\frac{3}{8}$	$\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{5}{8}$						
				$\frac{n+11}{32}$	$\frac{n-21}{32}$	$\frac{n+19}{32}$	$\frac{n-13}{32}$	$\frac{n+3}{32}$	$\frac{n-29}{32}$	$\frac{n+27}{32}$	$\frac{n-5}{32}$
32	5	$\frac{5}{8}$	$\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{3}{8}$						
				$\frac{n+22}{32}$	$\frac{n-10}{32}$	$\frac{n+30}{32}$	$\frac{n-2}{32}$	$\frac{n+6}{32}$	$\frac{n-26}{32}$	$\frac{n-18}{32}$	$\frac{n+14}{32}$
32	7	$\frac{7}{4}$	$\frac{7}{8}$	$\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{1}{8}$					
				$\frac{n+26}{32}$	$\frac{n+58}{32}$	$\frac{n+2}{32}$	$\frac{n+34}{32}$	$\frac{n-14}{32}$	$\frac{n+18}{32}$	$\frac{n-22}{32}$	$\frac{n+10}{32}$

Table 2: The correction terms of all possible $Y = \left(-1; \frac{1}{2}, \frac{1}{2}, \frac{m}{n}\right)$ listed by $p = 4m = |H_1(Y)|$ and $n \bmod m$.

and $p/q = 1$ with $g(K) = 1$.

Theorem 7 demands that the symmetrized Alexander polynomial of K be

$$\Delta_K(T) = T - 1 + T^{-1}.$$

By Theorem 8, we know that K is fibered. Therefore, thanks to the classical results found in, for example, Burde and Zieschang [BZ85], K is the right- or left-handed trefoil. By Theorem 4, only 1-surgery on the right-handed trefoil (or -1-surgery on the left) is elliptic. $\square$

Proof of Theorems 2, 3. The proofs are comparable. We will explicitly demonstrate the proof for $p = 4$.

As above, assume surgery coefficient is $4/q \geq 0$ with $(4, q) = 1$.

By Theorem 4, Y is either a lens space or

$$Y = \left(-1; \frac{1}{2}, \frac{1}{2}, \frac{1}{n}\right),$$

where $(n, 4) = 1$.

All of these manifolds are L-spaces, so Theorem 9 restricts q:

$$\frac{4}{q} \geq 2g(K) - 1$$

and either $p/q = 4/3$ with $g(K) = 1$ or $p/q = 4$ with $g(K) = 1$ or 2.

If $p/q = 4/3$, Theorem 7 shows that the symmetrized Alexander polynomial of K is

$$\Delta_K(T) = T - 1 + T^{-1},$$

so K is a trefoil.

On the other hand, if $p/q = 4$ and $g(K) \leq 2$, the Alexander polynomial may be any one of

$$\Delta_1(T) = T - 1 + T^{-1} \qquad (2)$$

$$\Delta_2(T) = T^2 - 1 + T^{-2} \qquad (3)$$

$$\Delta_3(T) = T^2 - T + 1 - T^{-1} + T^{-2}. \qquad (4)$$

To narrow this down further, calculate the corresponding correction terms as in Theorem 10:

$$\begin{aligned} d(L_{4,1},i) &= -\left(\frac{pq-(2i+1-p-q)^2}{4pq}\right) - d(L_{q,r},j) \\ &= -\left(\frac{4-(2i+1-4-1)^2}{16}\right) - d(L_{1,0},0) = \begin{cases} 3/4 & i=0 \\ 0 & i=\pm 1 \\ -1/4 & i=2 \end{cases} \end{aligned}$$

$$\begin{aligned} d(S^3_4(K),i) &= d(L_{4,1},i) - 2\sum_{j=1}^{\infty} j a_{j+c} \\ &= d(L_{4,1},i) + \left\{ \begin{array}{rrrl} \Delta_1 & \Delta_2 & \Delta_3 & \\ -2 & -4 & -2 & i=0 \\ 0 & -2 & -2 & i=\pm 1 \\ 0 & 0 & 0 & i=2 \end{array} \right. \\ &= \left\{ \begin{array}{rrrl} \Delta_1 & \Delta_2 & \Delta_3 & \\ -5/4 & -13/4 & -5/4 & i=0 \\ 0 & -2 & -2 & i=\pm 1 \\ -1/4 & -1/4 & -1/4 & i=2 \end{array} \right. \end{aligned}$$

Compare the correction terms calculated in Example 14. These correction terms match the 4-surgery correction terms only for $\Delta_1(T)$ when $n=3$, but $\Delta_1(T)$ is only the Alexander polynomial of the trefoil. Theorems 4 and 5 tell us that only one of the $\pm 4, \pm 4/3$- surgeries on the trefoil gives an elliptic Seifert fibered space:

$$\left(-1; \frac{1}{2}, \frac{1}{2}, \frac{1}{3}\right) = S^3_4(T_{3,2}) = -S^3_{-4}(T_{3,-2})$$

The cases of $\mathbb{Z}_2$ and $\mathbb{Z}_3$ are similar. For $\mathbb{Z}_2$, the surgery coefficient must be 2/1 with $g(K)=1$ and $\Delta_K(T) = T - 1 + T^{-1}$, and so K is the trefoil. For $\mathbb{Z}_3$, it must be 3/2 with $g(K)=1$ and K the trefoil or else 3 with $g(K) \leq 2$ and the Alexander polynomials from (4). As before, the only possible finite surgeries are again 2- or 3- surgery on the right-handed trefoil; the others are eliminated because the correction terms of the manifold in Table 2 do not match those of the possible finite surgeries in Table 3. $\square$

Δ_i	p	q	n											
Δ_1	4	1	-3	$-\frac{5}{4}$	0	$-\frac{1}{4}$								
Δ_1	8	1	3	$\frac{1}{4}$	$\frac{7}{8}$	$-\frac{1}{4}$	$-\frac{1}{8}$							
Δ_3	8	1	-5	$\frac{1}{4}$	$-\frac{9}{8}$	$-\frac{1}{4}$	$-\frac{1}{8}$							
Δ_3	12	1	5	$\frac{3}{4}$	$-\frac{1}{6}$	$\frac{13}{12}$	$\frac{1}{2}$	$\frac{1}{12}$	$-\frac{1}{6}$	$-\frac{1}{4}$				
Δ_4	12	1	-7	$-\frac{5}{4}$	$-\frac{1}{6}$	$-\frac{11}{12}$	$\frac{1}{2}$	$\frac{1}{12}$	$-\frac{1}{6}$	$-\frac{1}{4}$				
Δ_7	16	1	-1	$-\frac{1}{4}$	$\frac{13}{16}$	0	$-\frac{11}{16}$	$\frac{3}{4}$	$\frac{5}{16}$	0	$-\frac{3}{16}$	$-\frac{1}{4}$		
Δ_4	16	1	7	$-\frac{1}{4}$	$\frac{13}{16}$	0	$\frac{21}{16}$	$\frac{3}{4}$	$\frac{5}{16}$	0	$-\frac{3}{16}$	$-\frac{1}{4}$		
Δ_6	16	1	-9	$-\frac{1}{4}$	$-\frac{19}{16}$	0	$-\frac{11}{16}$	$\frac{3}{4}$	$\frac{5}{16}$	0	$-\frac{3}{16}$	$-\frac{1}{4}$		
Δ_{12}	20	1	1	$\frac{3}{4}$	$-\frac{1}{5}$	$-\frac{21}{20}$	$\frac{1}{5}$	$-\frac{9}{20}$	-1	$\frac{11}{20}$	$\frac{1}{5}$	$-\frac{1}{20}$	$-\frac{1}{5}$	$-\frac{1}{4}$
Δ_6	20	1	9	$\frac{3}{4}$	$-\frac{1}{5}$	$\frac{19}{20}$	$\frac{1}{5}$	$\frac{31}{20}$	1	$\frac{11}{20}$	$\frac{1}{5}$	$-\frac{1}{20}$	$-\frac{1}{5}$	$-\frac{1}{4}$
Δ_9	20	1	-11	$-\frac{5}{4}$	$-\frac{1}{5}$	$-\frac{21}{20}$	$\frac{1}{5}$	$-\frac{9}{20}$	1	$\frac{11}{20}$	$\frac{1}{5}$	$-\frac{1}{20}$	$-\frac{1}{5}$	$-\frac{1}{4}$
Δ_{12}	24	1	11	$-\frac{1}{4}$	$-\frac{29}{24}$	$-\frac{1}{12}$	$-\frac{7}{8}$	$\frac{5}{12}$	$-\frac{5}{24}$	$\frac{5}{4}$	$\frac{19}{24}$	$\frac{5}{12}$	$\frac{1}{8}$	$-\frac{1}{12}$
				$-\frac{5}{24}$	$-\frac{1}{4}$									
Δ_9	24	1	-13	$-\frac{1}{4}$	$\frac{19}{24}$	$-\frac{1}{12}$	$\frac{9}{8}$	$\frac{5}{12}$	$\frac{43}{24}$	$\frac{5}{4}$	$\frac{19}{24}$	$\frac{5}{12}$	$\frac{1}{8}$	$-\frac{1}{12}$
				$-\frac{5}{24}$	$-\frac{1}{4}$									
Δ_{16}	28	1	-5	$\frac{3}{4}$	$-\frac{3}{14}$	$\frac{25}{28}$	$\frac{1}{14}$	$-\frac{19}{28}$	$\frac{9}{14}$	$\frac{1}{28}$	$-\frac{1}{2}$	$\frac{29}{28}$	$\frac{9}{14}$	$\frac{9}{28}$
				$\frac{1}{14}$	$-\frac{3}{28}$	$-\frac{3}{14}$	$-\frac{1}{4}$							
Δ_{17}	28	1	9	$\frac{3}{4}$	$-\frac{3}{14}$	$-\frac{31}{28}$	$\frac{1}{14}$	$-\frac{19}{28}$	$\frac{9}{14}$	$\frac{1}{28}$	$-\frac{1}{2}$	$-\frac{27}{28}$	$\frac{9}{14}$	$\frac{9}{28}$
				$\frac{1}{14}$	$-\frac{3}{28}$	$-\frac{3}{14}$	$-\frac{1}{4}$							
Δ_{14}	28	1	13	$\frac{-5}{4}$	$-\frac{3}{14}$	$-\frac{31}{28}$	$\frac{1}{14}$	$-\frac{19}{28}$	$\frac{9}{14}$	$\frac{1}{28}$	$\frac{3}{2}$	$\frac{29}{28}$	$\frac{9}{14}$	$\frac{9}{28}$
				$\frac{1}{14}$	$-\frac{3}{28}$	$-\frac{3}{14}$	$-\frac{1}{4}$							
Δ_{12}	28	1	-15	$\frac{3}{4}$	$-\frac{3}{14}$	$\frac{25}{28}$	$\frac{1}{14}$	$\frac{37}{28}$	$\frac{9}{14}$	$\frac{57}{28}$	$\frac{3}{2}$	$\frac{29}{28}$	$\frac{9}{14}$	$\frac{9}{28}$
				$\frac{1}{14}$	$-\frac{3}{28}$	$-\frac{3}{14}$	$-\frac{1}{4}$							
$\Delta_{18,15}$	32	1	9	$\frac{7}{4}$	$\frac{25}{32}$	$-\frac{1}{8}$	$-\frac{31}{32}$	$\frac{1}{4}$	$\frac{49}{32}$	$\frac{7}{8}$	$\frac{9}{32}$	$-\frac{1}{4}$	$-\frac{23}{32}$	$\frac{7}{8}$
				$\frac{17}{32}$	$\frac{1}{4}$	$\frac{1}{32}$	$-\frac{1}{8}$	$-\frac{7}{32}$	$-\frac{1}{4}$					
Δ_{14}	32	1	-15	$-\frac{1}{4}$	$\frac{25}{32}$	$-\frac{1}{8}$	$\frac{33}{32}$	$\frac{1}{4}$	$\frac{49}{32}$	$\frac{7}{8}$	$\frac{73}{32}$	$\frac{7}{4}$	$\frac{41}{32}$	$\frac{7}{8}$
				$\frac{17}{32}$	$\frac{1}{4}$	$\frac{1}{32}$	$-\frac{1}{8}$	$-\frac{7}{32}$	$-\frac{1}{4}$					

Table 3: Correction terms for a p/q-surgery on a knot with Alexander polynomial Δ_i, assuming that it is an L-space surgery; also listed is the n for which some $(-1; \frac{1}{2}, \frac{1}{2}, \frac{m}{n})$ satisfies these correction terms.

Δ_i	a_0	a_1	a_2	a_3	a_4	a_5	a_6	a_7	a_8	a_9	a_{10}
Δ_1	-1	1									
Δ_2	-1	0	1								
Δ_3	1	-1	1								
Δ_4	-1	1	-1	1							
Δ_5	1	0	-1	1							
Δ_6	1	-1	1	-1	1						
Δ_7	-1	1	0	-1	1						
Δ_8	1	0	0	-1	1						
Δ_9	-1	1	-1	1	-1	1					
Δ_{10}	-1	1	0	0	-1	1					
Δ_{11}	1	0	0	0	-1	1					
Δ_{12}	1	-1	1	-1	1	-1	1				
Δ_{13}	1	0	-1	1	0	-1	1				
Δ_{14}	-1	1	-1	1	-1	1	-1	1			
Δ_{15}	1	0	-1	1	-1	1	-1	1			
Δ_{16}	1	-1	1	0	-1	1	0	-1	1		
Δ_{17}	1	0	-1	1	-1	1	0	0	-1	1	
Δ_{18}	1	0	0	-1	0	1	0	0	0	-1	1

Table 4: The Alexander polynomials used in Table 3.

In Table 1, we stated the results for the elliptic manifolds with $|H_1(Y)| \leq 10$ and for all other dihedral manifolds (which we find to be more interesting) with $|H_1(Y)| \leq 32$. For each choice of $p = |H_1(Y)|$, Theorem 4 gives a description like $(-1; \frac{1}{2}, \frac{1}{2}, \frac{p/4}{n})$ for all possible Y. The correction terms for each such manifold are listed in Table 2. On the other hand, assuming Y is a knot surgery, Theorem 9 restricts the surgery coefficients that can give Y, and Theorems 7 and 8 restrict the Alexander polynomial of the knot (of which there are 2^x where $x = \lfloor \frac{p}{2q} - \frac{1}{2} \rfloor$). Using Theorem 10, it is possible to calculate the correction terms of the resulting surgeries for each knot (assuming they are indeed L-spaces), which are shown in Table 3 with the corresponding Alexander polynomials listed in Table 4.

5 Conjectures

Torus knot surgeries are suspiciously common among the SFSs we calculated. It is tempting to conjecture:

Conjecture 15. *If an elliptic SFS (other than a lens space) is a knot surgery, then it is surgery on a torus knot [of form $T_{2,n}, T_{3,4}$, or $T_{3,5}$].*

Although perhaps it is safer to say:

Conjecture 16. *If an elliptic SFS (other than a lens space) is a knot surgery, then it is fibered and shares its Alexander polynomial with a $T_{2,n}, T_{3,4}$, or $T_{3,5}$ torus knot.*

Additionally, the correction terms of a plumbing diagram (when the diagram is the boundary of a negative definite 4-manifold) seem to be dependent only on $p = |H_1(Y)|$.It appears, however, that the numerators almost never match up for $q \neq 1$.

Proposition 17. *If $H_1(Y) = \mathbb{Z}_p$, then*

$$d(Y; \mathfrak{t}) \equiv \frac{c_1(\mathfrak{t})^2}{4} \bmod 1$$

Proof. This fact is obviously true from the relation $d(Y(G), \mathfrak{t}) = \max(k^2 + |G|)/4$ when G has at most two bad vertices, but it is true in general. (Thanks to Zoltán Szabó for explaining the reasoning for the general case.)

If $H_1(Y) = \mathbb{Z}_p$, then Y is obtained from an integral homology sphere Y' by p-surgery on a knot (find a simple closed curve K in Y that generates $H_1(Y)$ and do $1/p$-surgery on it to kill $H_1(Y; \mathbb{Q})$.) Also, Y' is obtained from S^3 by a finite series of ±1 surgeries (see, e.g., [Sav99]).

Consider the cobordism X_1 from Y to Y' which is Y union a 2-handle framed by $1/p$ along K and the cobordism X_2 from Y' to S^3 which is S^3 union some ± 1-framed 2-handles. A Spin^c structure on Y' extends to a Spin^c structure on X_2 when the map $H^2 X_2 \to H^2 Y'$ (induced by the boundary map in the long exact sequence) is surjective, which it is since $H^2 Y' = 0$. Similarly, a Spin^c structure on Y extends to one on X_1 since $H^2 X_1 \to H^2 Y$ is $H_2 X_1 \cong \mathbb{Z} \to H_1(Y) \cong \mathbb{Z}_p$ which takes the 2-handle to K.

$X = X_1 \cup X_2$ is a cobordism as well with induced map $f^+_{X,\mathfrak{s}} : CF^+(Y, \mathfrak{s}|_Y) \to CF^+(S^3, \mathfrak{s}|_{S^3})$. The relative $\mathbb{Z}$-grading on $HF^+(Y, \mathfrak{s}|_Y)$ therefore lifts to a $\mathbb{Q}$-grading consistent with S^3's grading, as in Equation **??**: $\mathrm{gr}(f^+_{X,\mathfrak{s}}(\xi)) = \mathrm{gr}(\xi) + (c_1(\mathfrak{t})^2 - 2\chi(X) - 3\sigma(X))/4 \equiv c_1(\mathfrak{t})^2/4 \bmod 1$. □

These $c_1(\mathfrak{t})^2$ also seem to display a few patterns: there always seem to be about m pairs of characteristic vectors whose squares are constant and thus give correction terms which are dependent upon n; these seem to come in pairs: $n + a/4m$, $n - b/4m$ for $a + b = 4m$. Generally, a pair occurs with one element in Spin^c structure $\pm i$ and the other in $n/2 \pm i$. Also, the other m pairs seem to be small numbers (independent of n) between 0 and 1. If $n = mk + p$ and $n' = (m + 1)k - p$, then n and n' have the same correction terms for the same Spin^c structures but with opposite signs.

Conjecture 18. *If $Y = \left(-1; \frac{1}{2}, \frac{1}{2}, \frac{m}{n}\right)$ with $n > m$ and $\mathfrak{t} \in Spin^c(Y)$ is torsion, then $c_1(\mathfrak{t})^2$ has denominator m (or a divisor of m).*

References

[Ber] John Berge, *Obtaining lens spaces by surgeries on knots*, unpublished.

[BH92] Steven A. Bleiler and Craig D. Hodgson, *Spherical space forms and Dehn surgery*, Knots 90 (Osaka, 1990), de Gruyter, Berlin, 1992, pp. 425–433. MR MR1177439 (93k:57008)

[BH96] ——, *Spherical space forms and Dehn filling*, Topology **35** (1996), no. 3, 809–833. MR MR1396779 (97f:57007)

[BZ85] Gerhard Burde and Heiner Zieschang, *Knots*, de Gruyter Studies in Mathematics, vol. 5, Walter de Gruyter & Co., Berlin, 1985. MR MR808776 (87b:57004)

[BZ96] S. Boyer and X. Zhang, *Finite Dehn surgery on knots*, J. Amer. Math. Soc. **9** (1996), no. 4, 1005–1050. MR 1333293 (97h:57013)

[BZ01] Steven Boyer and Xingru Zhang, *A proof of the finite filling conjecture*, J. Differential Geom. **59** (2001), no. 1, 87–176. MR 1909249 (2003k:57007)

[CGLS88] M. Culler, C. McA. Gordon, J. Luecke, and P. B. Shalen, *Correction to: "Dehn surgery on knots" [Ann. of Math. (2)* ***125*** *(1987), no. 2, 237–300; MR0881270 (88a:57026)]*, Ann. of Math. (2) **127** (1988), no. 3, 663. MR MR942524 (89c:57015)

[Dea03] John C. Dean, *Small Seifert-fibered Dehn surgery on hyperbolic knots*, Algebr. Geom. Topol. **3** (2003), 435–472 (electronic). MR MR1997325 (2004m:57009)

[DMM09] Arnaud Deruelle, Katura Miyazaki, and Kimihiko Motegi, *Networking Seifert surgeries on knots. II. The Berge's lens surgeries*, Topology Appl. **156** (2009), no. 6, 1083–1113. MR MR2493370

[Doi10] Margaret Doig, *Spherical seifert fibered spaces, knot surgeries, and heegaard floer homology*, Ph.D. thesis, Princeton University, 2010.

[EM02] Mario Eudave-Muñoz, *On hyperbolic knots with Seifert fibered Dehn surgeries*, Proceedings of the First Joint Japan-Mexico Meeting in Topology (Morelia, 1999), vol. 121 1-2, 2002, pp. 119–141. MR MR1903687 (2003c:57005)

[Gab87] David Gabai, *Foliations and the topology of* 3*-manifolds. III*, J. Differential Geom. **26** (1987), no. 3, 479–536. MR MR910018 (89a:57014b)

[GL89] C. McA. Gordon and J. Luecke, *Knots are determined by their complements*, Bull. Amer. Math. Soc. (N.S.) **20** (1989), no. 1, 83–87. MR MR972070 (90a:57006b)

[Gor83a] C. McA. Gordon, *Dehn surgery and satellite knots*, Trans. Amer. Math. Soc. **275** (1983), no. 2, 687–708. MR MR682725 (84d:57003)

[Gor83b] ______, *Dehn surgery and satellite knots*, Trans. Amer. Math. Soc. **275** (1983), no. 2, 687–708. MR 682725 (84d:57003)

[Gre] Joshua Greene, *The lens space realization problem*, unpublished.

[GS99] Robert E. Gompf and András I. Stipsicz, 4*-manifolds and Kirby calculus*, Graduate Studies in Mathematics, vol. 20, American Mathematical Society, Providence, RI, 1999. MR MR1707327 (2000h:57038)

[Juh06] András Juhász, *Holomorphic discs and sutured manifolds*, Algebr. Geom. Topol. **6** (2006), 1429–1457 (electronic). MR MR2253454 (2007g:57024)

[Kir78a] Rob Kirby, *Problems in low dimensional manifold theory*, Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 2, Proc. Sympos. Pure Math., XXXII, Amer. Math. Soc., Providence, R.I., 1978, pp. 273–312. MR 520548 (80g:57002)

[Kir78b] Robion Kirby, *A calculus for framed links in* S^3, Invent. Math. **45** (1978), no. 1, 35–56. MR MR0467753 (57 #7605)

[Lic62] W. B. R. Lickorish, *A representation of orientable combinatorial* 3*-manifolds*, Ann. of Math. (2) **76** (1962), 531–540. MR MR0151948 (27 #1929)

[MMM06] Thomas Mattman, Katura Miyazaki, and Kimihiko Motegi, *Seifert-fibered surgeries which do not arise from primitive/Seifert-fibered constructions*, Trans. Amer. Math. Soc. **358** (2006), no. 9, 4045–4055 (electronic). MR MR2219009 (2007h:57009)

[Mos71] Louise Moser, *Elementary surgery along a torus knot*, Pacific J. Math. **38** (1971), 737–745. MR MR0383406 (52 #4287)

[Ni09] Yi Ni, *Link Floer homology detects the Thurston norm*, Geom. Topol. **13** (2009), no. 5, 2991–3019. MR MR2546619

[OS] Peter S. Ozsváth and Zoltán Szabó, *Knot Floer homology and rational surgeries*, unpublished.

[OS03a] Peter Ozsváth and Zoltán Szabó, *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, Adv. Math. **173** (2003), no. 2, 179–261. MR MR1957829 (2003m:57066)

[OS03b] ———, *On the Floer homology of plumbed three-manifolds*, Geom. Topol. **7** (2003), 185–224 (electronic). MR MR1988284 (2004h:57039)

[OS04a] ———, *Holomorphic disks and genus bounds*, Geom. Topol. **8** (2004), 311–334 (electronic). MR MR2023281 (2004m:57024)

[OS04b] ———, *Holomorphic disks and three-manifold invariants: properties and applications*, Ann. of Math. (2) **159** (2004), no. 3, 1159–1245. MR MR2113020 (2006b:57017)

[OS04c] ———, *Holomorphic disks and topological invariants for closed three-manifolds*, Ann. of Math. (2) **159** (2004), no. 3, 1027–1158. MR MR2113019 (2006b:57016)

[OS04d] ———, *On heegaard floer homology and seifert fibered surgeries*, Geom. Topol. Mono. **7: Proceedings of the Casson Fest** (2004), 181–203 (electronic).

[OS05] ———, *On knot Floer homology and lens space surgeries*, Topology **44** (2005), no. 6, 1281–1300. MR MR2168576 (2006f:57034)

[OS06] ———, *Holomorphic triangles and invariants for smooth four-manifolds*, Adv. Math. **202** (2006), no. 2, 326–400. MR MR2222356 (2007i:57029)

[OS08] Peter S. Ozsváth and Zoltán Szabó, *Knot Floer homology and integer surgeries*, Algebr. Geom. Topol. **8** (2008), no. 1, 101–153. MR MR2377279 (2008m:57075)

[Pera] Grisha Perelman, *The entropy formula for the ricci flow and its geometric applications*, unpublished.

[Perb] ———, *Finite extinction time for the solutions to the ricci flow on certain three-manifolds*, unpublished.

[Perc] ———, *Ricci flow with surgery on three-manifolds*, unpublished.

[Sav99] Nikolai Saveliev, *Lectures on the topology of* 3*-manifolds*, de Gruyter Textbook, Walter de Gruyter & Co., Berlin, 1999, An introduction to the Casson invariant. MR MR1712769 (2001h:57024)

[Sco83] Peter Scott, *The geometries of* 3*-manifolds*, Bull. London Math. Soc. **15** (1983), no. 5, 401–487. MR MR705527 (84m:57009)

[Sei33] H. Seifert, *Topologie Dreidimensionaler Gefaserter Räume*, Acta Math. **60** (1933), no. 1, 147–238. MR MR1555366

[ST80] Herbert Seifert and William Threlfall, *Seifert and Threlfall: a textbook of topology*, Pure and Applied Mathematics, vol. 89, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New

York, 1980, Translated from the German edition of 1934 by Michael A. Goldman, With a preface by Joan S. Birman, With “Topology of 3-dimensional fibered spaces” by Seifert, Translated from the German by Wolfgang Heil. MR MR575168 (82b:55001)

[Thu82] William P. Thurston, *Three-dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. (N.S.) **6** (1982), no. 3, 357–381. MR 648524 (83h:57019)

[Wal63] Andrew H. Wallace, *Modifications and cobounding manifolds. IV*, J. Math. Mech. **12** (1963), 445–484. MR MR0154297 (27 #4246)

[Zha93] Xingru Zhang, *On property I for knots in S^3*, Trans. Amer. Math. Soc. **339** (1993), no. 2, 643–657. MR 1154545 (93m:57011)